

## Reaction of Anti-Tubercular Isonicotinic Acid Hydrazide Drug with Chlorides of Niobium (V) and Tantalum (V)

R.N. Pandey<sup>1,\*</sup>, Pramila Sharma<sup>2</sup> and Sachchidanand Choudhary<sup>1</sup>

<sup>1</sup>P.G. Centre of Chemistry (M.U.), College of Commerce, Patna- 800020 (India)

<sup>2</sup>Department of Chemistry, Ganga Devi Mahila Mahavidyalaya, Patna- 800020

E-mail: rameshwarnath.pandey@yahoo.com

Article History:

Received: 15 July 2013

Accepted: 12 December 2013

### ABSTRACT

The reaction products of niobium (V) and tantalum (V) Chlorides with anti-Tubercular isonicotinic acid hydrazide drug are isolated and their structures are deduced using elemental analysis, conductivity measurements, magnetic susceptibility, IR and UV-vis Spectral studies. All metal derivatives were seven-coordinated having Pentagonal bipyramidal structure and isonicotinic acid hydrazide drug acts as bidentate NO donor. Metal-ligand vibrations in far-IR Spectra of complexes are analyzed and tentatively assigned.

**Keywords:** Niobium, Tantalum, Isonicotinic acid, Isoniazide.

©2013 ijCEPr. All rights reserved

### INTRODUCTION

A part of our studies on reaction between anti-Tubercular isonicotinic acid hydrazide drug with low-valent [1] and high valent [2] metal ions. We have studied reactions involving the niobium (V) and Tantalum (V) Chlorides. The formation of O-abstraction products by  $MCl_5$  ( $M=Nb/Ta$ ) above room temperature was also observed leading to the formation of solid products. The different solid products in different reaction conditions were isolated with coordination number seven having Pentagonal bipyramidal structure. The ligand-bridged and chloro-bridged structures were observed and deduced using various physico-chemical measurements.

### MATERIALS AND METHODS

All the reagents were AR grade or CP grade.  $MX_5$  ( $M = Nb/Ta$ ;  $X=Cl$ ) (Aldrich) and isonicotinic acid hydrazide (E.Merch) were commercial products of the highest purity available. All complexes were prepared using a general method taking  $MX_5$  and ligand in different molar ratios in  $CCl_4:CH_3OH$  (1:1) solvents adopting our previous method [3].

Elemental analysis was performed by micro-analysis division, RSIC, CDRI, Lucknow. IR Spectra of ligands and complexes were recorded on a Perkin-Elmer model- 577 Spectrophotometer in the range of  $4000-200\text{ cm}^{-1}$  as KBr pellets. The electronic spectra were recorded with Ziess (Jena) Model of automatic recording system. Magnetic measurements were made on Gouy balance using Hg  $[CO(SCN)_4]$  as calibrant. The molar conductance of complexes ( $10^{-3}M$ ) were measured in DMF using Wiss- Wekstatter Weitheim obb type LBR conductivity meter. Niobium and tantalum was determined gravimetrically as pentaoxide and chloride as silver chloride.

### RESULTS AND DISCUSSION

The reaction between  $MCl_5$  and isonicotinic acid hydrazide in  $CCl_4 : MeOH$  (1:1) yielded various products given in table 1. The reaction between them is exothermic and the nature of products formed was dependent on the stoichiometry of reacting species, pH value of reaction mixture and temperature. At room temperature no evidences for the formation of O-abstraction products were found. In all the reactions of  $MCl_5$  an addition compound was probably first formed after which  $HCl(g)$  was liberated. All niobium (V) complexes showed slight paramagnetism between 0.37 - .53 BM which is in agreement with results reported earlier [4]. The products did not melt but were decomposed to blue black mass in all cases in the temperature range  $235-250^\circ C$  due to reduction of Nb (V) to blue Nb (III) and Ta (V) to black Ta (III) by ligand [5]. The molar conductance in DMF ( $10^{-3}M$ ) were observed between  $20-32. \text{ ohm}^{-1}\text{cm}^2\text{mol}^{-1}$  suggested their non electrolyte nature [6]. However, molar conductance of  $[TaL_3Cl] Cl$  (S.No. 4) was  $89.72 \text{ ohm}^{-1}\text{cm}^2\text{mol}^{-1}$  indicating as 1:1 electrolytes [7]. The similar orientation of ligand about the metal ion was indicated having same formula type.

Electronic spectra of complexes exhibit strong band in the region 28170-27625  $\text{cm}^{-1}$  due to charge transfer besides the ligand bands. However, no absorption is observed above 25000  $\text{cm}^{-1}$  indicating  $d^0$ - configuration of Nb (V) and Ta (V) complexes.

### IR Spectra


The characteristic IR bands at 3400  $\text{cm}^{-1}$ , 3310  $\text{cm}^{-1}$ , 3210  $\text{cm}^{-1}$ , 1680  $\text{cm}^{-1}$ , 1560  $\text{cm}^{-1}$  in the spectrum of ligand are assigned to  $\nu\text{NH}_2$ ,  $\nu_{\text{asym}}(\text{NH})$ ,  $\nu_{\text{sym}}(\text{NH})$ , amide band I and amide band II respectively. The  $\nu\text{NH}_2$  of free ligand red shift to lower frequency 100  $\text{cm}^{-1}$  on coordination indicating bonding of ligand through amino nitrogen of hydrazine residue[1, 2]. The red shift (25-30  $\text{cm}^{-1}$ ) of band observed at 1640  $\text{cm}^{-1}$  due to  $\text{NH}_2$  deformation associated with OCN bending on complexation also support the formation of M-N bond. The pronounced shifting of  $\nu\text{C-N} + \nu\text{N-H}$  frequency by 30-35  $\text{cm}^{-1}$  towards lower frequency in complexes as compared to free ligand also supports involvement of amino nitrogen of the ligand in coordination. New bands at 503  $\text{cm}^{-1}$ , 524  $\text{cm}^{-1}$  & 480  $\text{cm}^{-1}$   $\nu\text{Nb-N}$  and at 546  $\text{cm}^{-1}$ , 510  $\text{cm}^{-1}$  and 450  $\text{cm}^{-1}$   $\nu\text{Ta-N}$  in far- IR Spectra of complexes also support formation of metal-N bond[8].

Table-1: Analytical and physical data of complexes

S. No.	Compound (pH, color)	% analysis : Calcd/ (Found)					$\Delta m$ $\text{ohm}^{-1}$ $\text{cm}^2$ $\text{mol}^{-1}$
		C	H	N	Cl	M	
1	[NbOL <sub>3</sub> ] (7, Faint Pink)	56.69 (56.82)	4.72 (4.83)	33.07 (33.32)	-	24.41 (24.50)	31.48
2	[TaOL <sub>3</sub> ].2CH <sub>3</sub> OH (7, Brown)	35.87 (36.21)	3.88 (3.43)	18.40 (19.90)	-	27.05 (27.11)	32.43
3	[NbL <sub>3</sub> Cl]Cl (7.5, Pale yellow)	37.76 (37.72)	3.14 (3.66)	22.02 (22.01)	12.41 (12.50)	16.25 (16.12)	73.60
4	[TaL <sub>3</sub> Cl]Cl (7.5, Yellow)	32.72 (32.83)	2.72 (2.71)	19.09 (19.32)	10.75 (10.80)	27.42 (27.50)	89.72
5	[Ta <sub>2</sub> (LH)(H <sub>2</sub> O)Cl <sub>10</sub> ] (6, Orange)	7.62 (7.37)	1.80 (1.67)	2.40 (2.38)	37.60 (36.82)	38.34 (38.50)	23.02
6	[Ta <sub>2</sub> (LH)(NCS) <sub>2</sub> Cl <sub>8</sub> (CH <sub>3</sub> OH) <sub>2</sub> ] (6, Yellow)	12.46 (12.26)	1.55 (1.77)	7.26 (7.54)	29.49 (29.50)	37.59 (37.62)	20.3
7	[Ta <sub>2</sub> (LH)Cl <sub>10</sub> (CH <sub>3</sub> OH) <sub>2</sub> ] (6, Brown)	10.45 (10.42)	1.63 (1.56)	4.57 (4.62)	38.67 (38.71)	39.43 (39.50)	21.20
8	[Nb <sub>2</sub> (LH)(H <sub>2</sub> O) <sub>2</sub> Cl <sub>10</sub> ] (6, Orange Yellow)	10.08 (10.11)	1.54 (1.62)	5.88 (5.87)	49.71 (49.75)	26.05 (26.11)	22.32

(Where, LH = C<sub>6</sub>H<sub>7</sub>N<sub>3</sub>O)


The  $\nu\text{CO}$  of the ligand observed at 1680  $\text{cm}^{-1}$  as strong band red shift to lower frequency (35-40  $\text{cm}^{-1}$ ) on complexation indicating bonding through carbonyl oxygen atom which decreases CO bond order and increase in CN bond order substantiated by splitting of band at 1255  $\text{cm}^{-1}$  into two component (1195-1210 $\text{cm}^{-1}$ ) and (1380-1390 $\text{cm}^{-1}$ ) in the spectra of complexes (S. No. 5, 6, 7 & 8). The formation of M-O is indicated by the replacement of hydrogen atom of OH group of the enolic form (amidol system) of the ligand in other complexes (S. No. 1, 2, 3, 4).



The  $\nu_{\text{C-N}}$  observed at  $1490\text{ cm}^{-1}$  in free ligand shifted to  $1500\text{-}1515\text{ cm}^{-1}$  on complexation which suggest enol form seemed more prominent at high pH values where replacement of hydrogen atom or deprotonation appeared easier. New non-ligand bands around  $370\text{-}415\text{ cm}^{-1}$  supports the form of Metal-O bond [9].

The  $\nu_{\text{CN}}$ ,  $\nu_{\text{CS}}$  and  $\delta_{\text{NCS}}$  frequencies observed at  $2045\text{ cm}^{-1}$ ,  $785\text{ cm}^{-1}$  and  $490\text{ cm}^{-1}$  confirms the N-bonded isothiocyanate ion [10] in corresponding complex (S. No. 6). The presence of strong absorption in the  $850\text{-}950\text{ cm}^{-1}$  region also confirms the existence of  $\text{M=O}$  species in OXO- complexes (S.No. 1 and 2) [11]. New bands between  $340\text{-}345\text{ cm}^{-1}$  in niobium (V) complexes and at  $460\text{-}465\text{ cm}^{-1}$  in tantalum (V) complexes in far IR Spectra are assigned due to metal-Cl stretching mode [12]. Such bands are not present in OXO-complexes (S.No. 1 and 2). The bridging chloride group vibration in complexes is generally observed around  $200\text{ cm}^{-1}$  [13]. The new bands at  $250$  and  $283\text{ cm}^{-1}$  in far IR spectra does not ruled out the possibility of bridging chlorido group. The decrease of  $\nu_{\text{C-O}}$  (alcoholic) stretch of MeOH from  $1034\text{ cm}^{-1}$  to lower energy by  $45\text{-}70\text{ cm}^{-1}$  in complexes (S. No.6 and 7) indicate the involvement of the O atom of MeOH towards coordination[14].

Thus, on the basis of analytical data (Table-1), valence requirements and the infrared spectral studies, it is proposed that the ligand behaves as bidentate and structure of complexes may be deduced as:



## REFERENCES

1. Pandey R.N., Kumar Sunil, Kumar Shashikant, Choudhary Rita Roy and Sahay A.N., Indian J. Chem., **32A** (1993)987.
2. Pandey R.N., Singh Sarita and Sinha A.K., Asian J. Chem., **6(2)** (1994)252.
3. Pandey R.N. and Kumar S.S., J. Ultra Chem., **8(2)** (2012)135.
4. Kepert D.L. and Nyholm R.S., J. Chem. Soc., **Part II** (1965) 2871.
5. Cotton F.A. and Wilkinson G., Advanced Inorganic Chemistry, Inc Publisher, N.Y. P. ,(1976) 939.
6. Chabanur H.S., Revankar V.K. and Mahale V.B., Synth. React. Inorg. Met. Org. Chem., **31(2)** (2001) 339.
7. Geary W.J., Coord. Chem. Rev., 7 (1971)81.

8. Sharma R.K., Singh R.V. and Tandon J.P., J. Inorg. Nucl. Chem., **42** (1980)463.
9. Chabanur H.S., Revankar V.K. and Mahate V.B., Synth. React. Inorg. Met. Org. Chem., **31(2)** (2001) 339.
10. Burmister J.L., Coord. Chem. Rev., **1** (1966)205; **3** (1968) 225.
11. Heckley P.R. and Holah D.G., Can. J. Chem., **49** (1971)1151.
12. Gudasi K.B., Maravalli P.B. and Goudar T.R., J. Serb. Chem. Soc., **70(4)**(2005) 643.
13. Goldstein M. and Unsworth W.D., Inorg. Nucl. Chem. Lett., **6**(1970) 25.
14. Kumar D., Syamal A., Kumar A., Gupta P.K. and Dass D., J. Indian Chem. Soc., **87** (2010) 417.

[ijCEPr-266/2013]

**Water: Research & Development****[Water R&D]**

www.waterrnd.com

ISSN: 2249-2003

[Abstracted in : Chemical Abstracts Service, USA and CAB(I) , UK]

**WaterR&D** is an international Research Journal, dedicated to 'Water'. It is a truly interdisciplinary journal on water science and technology. It'll showcase the latest research related to Water in the field of chemistry, physics, biology, agricultural, food, pharmaceutical science, and environmental, oceanographic, and atmospheric science. It includes publication of reviews, regular research papers, case studies, communications and short notes.

*Manuscript Categories: Full-length paper, Review Articles, Short/Rapid Communications.*

Manuscripts should be addressed to:

**E-mail: waterrd@gmail.com**

*International Journal of***Chemical, Environmental and Pharmaceutical Research**

www.ijcepr.com; www.ijcepr.in

ISSN: 2229-3892(Print); ISSN: 2229-5283(Online)

[Abstracted in : Chemical Abstracts Service , American Chemical Society, USA and CAB(I) , UK]

**ijCEPr** widely covers all fields of **Chemical, Environmental and Pharmaceutical Research.**

*Manuscript Categories: Full-length paper, Review Articles, Short/Rapid Communications.*

Manuscripts should be addressed to:

**E-mail: ijcepr@gmail.com**

**Adopt GREEN CHEMISTRY**

Save Our Planet.

We publish papers of Green Chemistry on priority.

If you think that you may be a potential reviewer in field of your interest, write us at [rasayanjournal@gmail.com](mailto:rasayanjournal@gmail.com) with your detailed resume and recent color photograph.